

# **SOME ASPECTS OF THE SLEEP OF LACTATING RAT DAMS**

Lúcia Regina Machado da Rocha<sup>1</sup> and Katsumasa Hoshino<sup>2</sup>

<sup>1</sup>Departamento de Fisiologia, Instituto de Biociências, Universidade Estadual Paulista, Campus Botucatu, São Paulo, Brasil

<sup>2</sup>Departamento de Ciências Biológicas, Faculdade de Ciências, Universidade Estadual Paulista, Campus Bauru, São Paulo, Brasil

Running title: Sleep of lactating rat dams

Corresponding author

L.R.M. Rocha - Depto. de Fisiologia - Instituto de Biociências - UNESP, 18618-000 BOTUCATU, S.P., Brasil.

**Acknowledgements:** Work supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP)

## ABSTRACT

The present study describes some aspects observed in the sleep of lactating rat dams (LRD) since literature on the subject is scanty. Electrographic recordings were made from 10:00h to 17:00h in nine rat dams, in different days after delivery (2<sup>nd</sup> to 20<sup>th</sup> day). Normal estrous cycling female rats, with previous motherhood experience, were used as controls. Four adult males were also used for comparisons. Sleep-wakefulness was quantified and expressed as percentages of the total recording time and significance assessed by ANOVA, adopting significance at the 0.05 level. Total wakefulness was  $43.4 \pm 2.6\%$  in control females,  $44.3 \pm 7.6\%$  in the males and  $55.1 \pm 4.2\%$  in LRD living with their 8 pups (difference rejected at the limit,  $p = 0.0547$ ). The mean number of awakening episodes longer than 30 minutes was significantly greater in LRD. Synchronized wave sleep amounted to  $46.6 \pm 2.1\%$ ,  $46.1 \pm 7.6\%$  and  $41.9 \pm 4.0\%$ , respectively in control females, males and LRD. Desynchronized wave sleep (REM-sleep) was significantly reduced in LRD ( $3.0 \pm 0.4\%$ ) compared to the values found in control females ( $10.0 \pm 1.0\%$ ) and males ( $9.4 \pm 0.6\%$ ). The average amounts of REM-sleep/hour along the recording period revealed to be constant and significantly lower in the lactating rat dams ( $3.0 \pm 1.3\%$ ) comparatively to control females ( $11.6 \pm 3.8\%$ ). An average of  $1.2 \pm 0.5$  REM-sleep episodes/hour was also significantly lower in lactating females than in control females ( $3.2 \pm 1.5$ ). The results obtained suggest an adaptive reorganization of the sleep architecture in the female rat during the nursing period.

**Key words:** sleep; nursing rats; lactation.

---

## INTRODUCTION

The rat is undoubtedly the animal most used in experimental sleep research and data on the subject is voluminous. Despite this, sleep studies using electrographic tools in order to quantify sleep in lactating rat dams (LRD) are scanty in the literature. Such scantiness was also recognized in humans by Karacan and colleagues (1968). In spite of the recognized occurrence of behavioral disorders in this period of women's lives, very few sleep studies were performed until now, except for those devoted to the dependence of milk ejection to bouts of synchronized sleep (SS), reported by Voloschin & Tramezzani (1979; 1984).

The present study aims to report some data on the sleep of LRD since they seem to represent important adaptations for this period of the female's life.

## METHODS

The sleep-wakefulness parameters were determined in 9 lactating rat dams, living in their home-cages with 8 pups each. Nine females displaying normal estrous cycles but having previous motherhood experience and 4 males were used as controls. All of the animals were 5-6 months old and were provided by the Central Vivarium of Universidade Estadual Paulista,

Botucatu, from a strain of albino rats (originally from the Wistar stock) reared for more than 20 years. Chronic electrodes were surgically implanted in all animals for electrocorticographic (ECoG), electromyographic (EMG) and electroculographic (EOG) recordings for determination of sleep-wakefulness parameters. Such determination was made from recordings obtained on days 2, 6, 9, 10, 11, 15, 16 19 and 20 of lactation (using only the first recording made in each animal). Four control female rats were in diestrous, 2 in proestrous and 3 in estrous phase on the morning of the electrographic recording day.

Electrodes for ECoG, EMG and EOG were chronically implanted in the rats under pentobarbital (40mg/kg) and local (Xylocain 2%, with vasoconstrictor) anesthesia. All recommendations for animal use in experimental research were followed. All procedures for surgical implantation of electrodes and recordings followed the descriptions made in previous publications (Timo-laria et al. 1970; Hoshino & Tolo, 1995). After surgery, the animals were moved to their plastic individual home cages (380x400x160mm), provided with commercial food pellets, potable water and wood shavings. A period of at least seven days for post-surgery recovery was allowed before recordings, maintaining the cages in a controlled 12:12h light-dark cycle (light on at 07:00h) and temperature room (24-25°C).

At 08:00h, the animal was moved in its home cage to a contiguous room and kept inside a Faraday's cage on the day of electrographic recording. Room temperature was kept constant (24-25°C) and a paper cardboard fixed on the wall of the Faraday's cage in order to prevent the direct incidence of the bulb lamp's light on the nest. Electrodes were connected to a Beckman polygraph and recording started at 10:00h and ended at 17:00h. Recordings were made always with paper speed at 5 or 10mm/second.

Two independent scorers, using 2-s epochs, visually analyzed recordings as awakening, SS or REM-sleep. The criteria used for the scoring of each state were similar to those adopted and described in previous quantitative studies (Timo-laria et al., 1970; Schmidek et al., 1972). A third experienced judge solved the cases of discordant scoring segments. All results were expressed as percentages of total recording time (TRT). Analysis of variance (ANOVA) (Statistica Software) was used to evaluate the existence of significant differences among groups and, when necessary, Tukey's honest significant difference (HSD) test was used as the *post-hoc* test. Significance level was adopted at  $p < 0.05$ .

## RESULTS

The prominent behavior of LRD was their permanence in the nest, where sleep, lactation and pup cleansing were displayed in an alternating fashion. The remaining part of

the time corresponded to brief episodes of exploratory activity around the cage, nest repair, eating, drinking, sporadic pup retrieval and sleep episodes outside the nest. These behaviors waned by the time of initial offspring's trials to eat food pellets by the 15<sup>th</sup> day, when adulthood pattern of locomotion was observed. The mean amount of time spent awake by LRD was  $55.1 \pm 4.2\%$  of the TRT. The greatest amounts were observed on the 6<sup>th</sup> and 11<sup>th</sup> lactation days (75.8 and 72.6%, respectively), whereas the lowest values were observed on the 10<sup>th</sup> and 18<sup>th</sup> days (44.5 and 38.6%, respectively). The statistical significance of the mean presented by the LRD group, comparatively to controls, was rejected at the limit (ANOVA,  $F_{(2,19)} = 3.397$ ,  $p=0.0547$ ), as shown in Table 1. The increased time spent awake by LRD was determined mainly by frequent pups' demand for suckling. They awoke periodically and synchronously, becoming agitated and presenting uncoordinated movements and strident vocalization while nipple-searching behavior was evident. Although the offspring slept approximately in a synchronized way, many intense phasic muscular jerks of sleep displayed by the newborns disturbed the lactating dam. In such cases, dams changed posture in order to facilitate nipple access or to attend other demands. Such facts increased the number of awakenings longer than 30 minutes in LRD (mean of  $1.37 \pm 0.31$  episodes/hour) that was significantly greater ( $F_{(2,19)} = 10.16$ ,  $p<0.05$ ) than those observed in control females ( $0.84 \pm 0.27$  episodes/hour). After eyes opening by the 11<sup>th</sup>-12<sup>th</sup> day of age, lactation demand became progressively silent; however, nipple-searching movements became more vigorous, which also awakened the dams. LRD progressively refused lactation when pups started to try solid food ingestion. However, dams continued to sleep on an overcrowded environment.

SS amounted to  $41.9 \pm 4.0\%$  of the TRT in the LRD. This value was not significantly different from that found in control groups (Table 1). The amount of REM-sleep in LRD never surpassed 5% of the TRT and the group reached an average of  $3.0 \pm 0.4\%$ . The amount presented by the LRD group was significantly lower ( $F_{(2,19)} = 24.948$ ,  $p<0.05$ ) than the means of control females (Tukey HSD test,  $p<0.05$ ) and males ( $p<0.05$ ), as presented in Table 1. The mean amounts of REM-sleep/hour, throughout the recording period, revealed to be steadily and significantly lower in lactating rat dams ( $3.0 \pm 1.3\%$ ) comparatively to control females ( $11.6 \pm 3.8\%$ ) ( $F_{(1,12)} = 32.770$ ,  $p<0.05$ ). The number of REM-sleep episodes presented by LRD ranged from 3 to 17, showing relatively higher values on the last days of lactation (mean 12.6 episodes) compared to the initial days (mean 6.6 episodes). Control female rats displayed about 20 episodes each during the similar 7 hours of recordings. The mean number of  $1.2 \pm 0.5$  episodes of REM-sleep/hour was significantly lower in LRD than the that observed in control

females ( $3.1 \pm 1.5$ ) ( $F_{(1,12)} = 9.434$ ,  $p < 0.05$ ). All longer episodes of REM-sleep observed in lactating dams were manifested outside the nest. In such case, they laid down alone and stretched to full length on the wooden shavings, distant from the pups.

**Table 1** - Parameters of diurnal sleep-wakefulness cycles in lactating rat dams and in control animals. Values are expressed as mean percentages ( $\pm$  s.d.) of the total recording time (from 10:00h to 17:00h). W = wakefulness; SS = synchronized sleep.

Parameter	Lactating mothers (n=9)	Cycling females (n=9)	Males (n=4)
Waking	$55.1 \pm 4.2^A$	$43.4 \pm 2.6$	$44.3 \pm 7.6$
SS	$41.9 \pm 4.0$	$46.6 \pm 2.1$	$46.1 \pm 7.6$
REM-sleep	$3.0 \pm 0.4^*$	$10.0 \pm 1.0$	$9.4 \pm 0.6$
Mean number of episodes per hour:			
a) Waking longer than 30 min	$1.4 \pm 0.3^*$	$0.8 \pm 0.3$	nc
b) REM-sleep	$1.2 \pm 0.5^*$	$3.1 \pm 1.5$	nc

\*- Significant difference (ANOVA plus Tukey's test,  $p < 0.05$ ); nc = non computed;

<sup>A</sup>- ( $p = 0.0547$ )

## DISCUSSION

The results obtained in the present study seem to disclose two important and interrelated changes in the sleep-wakefulness cycles of LRD. First, they displayed more awakening episodes longer than 30 min than their controls; second, they manifested lower amounts of REM-sleep.

The increase in the number of longer awakening episodes observed in LRD is similar to the ones observed in human mothers during the first month of lactation, as revealed by actigraphic and sleep-log monitoring techniques (Shinkoda, Matsumoto & Park 1999) or in the first two postpartum weeks as determined by EEG monitoring (Karacan et al., 1968). The

increase in wakefulness episodes showed by LRD does not seem to be a secondary consequence of the overcrowded condition inside the nest. Overcrowding was also reported to induce an increase in the number of wakefulness episodes (Sandrin, 1996); however, in such case, animals resume sleep. Awakening episodes in LRD are frequently devoted to several kinds of maternal care, and they may last more than 30 minutes, as observed in the present study. As constant maternal care is essential for the newborns, it may be thought the increase in awakening episodes as being an important adaptation for the lactation period.

Preservation of SS amounts in LRD seems to be also a significant adaptation for the nursing period. Milk-ejection reflex depends on previous bout of SS, being sleep probably induced by a rise in prolactin release (Voloschin & Tamezzaini, 1979; Voloschin & Tamezzaini, 1984; Lincoln et al., 1980). If one considers that increased amounts of wakefulness and preservation of SS are important adaptive manifestations during lactation, the reduced amount of REM-sleep in LRD, observed in the present study, may be taken as a mere remainder time of the recording period. This does not seem to be the case. Neither the case to attribute such fact to the impact of delivery and concomitant psychological changes as stated by Karacan and colleagues (1968) for the marked suppression of REM-sleep that occurs in the first postpartum day in humans. This REM-sleep change in humans is an acute suppression, therefore, not similar to that observed in LRD in which the state of sleep is manifested chronically and in reduced amounts during the lactation period.

Sleep in normal male adult rats is manifested mainly during the diurnal phase of the circadian cycle. In females the circadian distribution also exists, with a small variation in the REM-sleep amount associated to a change in the estrous cycle phase (Yamaoka, 1978; Yamaoka, 1980; Kimura et al., 1996). A regular daily amount of REM-sleep is necessary, and a need to compensate for its loss develops after a total or partial deprivation period (Dement, 1960). Maternal care is required by rat pups during the entire circadian cycle and this seems to impose a rearrangement on the mother's sleep-wakefulness architecture during the lactation period. Such reorganization explains and allows considering all results obtained in the present study as parts of the same adaptive mechanism. The need for a daily amount of REM-sleep seems to shift part of its diurnal manifestation to the nocturnal period, allowing the fulfillment of the need to spent more time awake and to preserve diurnal amounts of SS. In addition, SS needed for the milk-ejection reflex seems granted at night, since the lack of REM-sleep induces somnolence and promotes also an increase in SS amount during compensation. This parallel increase in SS amounts for REM-sleep compensation is a well-known fact (Jouvet, 1967). Although attractive, our few trials to detect signs of REM-sleep

compensatory rebound in LRD were considered negative since no significant changes were observed when they were allowed to sleep alone after pups' removal.

REM-sleep episodes manifested by LRD outside the nest seems an important topic among the behavioral data obtained in the present study. REM-sleep manifestation in stretched posture has a temperature-dissipating role (Schmidek et al., 1972) and such behavior in LRD suggests an important involvement of thermoregulatory mechanisms in their sleep organization. Body temperature in LRD is relatively higher (León et al., 1978; Bates et al., 1985) and this is important for pups' development (Adels & Leon, 1986). Prolonged maintenance of elevated back-posture on the nest overcrowded by pups seems the cause of overheating. Such overheating is possible since thermoregulatory capacity in LRD is lower (Knecht et al., 1980). The elevated basal temperature of LRD seems to be determined by the higher immune activity occurring in the LRD organism. Many agents of the immune system induce hyperthermia and promotes SS, and may reduce REM-sleep (Johansen et al., 1990; Krueger, 1994; Moldofsky, 1995). Sleep reduction may contribute also for the hyperthermia since its deprivation promotes an increase in body temperature (Landis et al., 1992; Seabra & Tufik, 1993; Hoshino, 1996). These properties of immune agents suggest their possible involvement in the sleep architecture reorganization observed in LRD.

It is important to note that our more recent studies indicate that the strain of Wistar rats used in the present study has some particular characteristics (Leneberg-Hoshino, 1999), like increased level of anxiety (dePaula, 1999) and is more susceptible to anxiety disorder-like manifestations when submitted to REM-sleep deprivation (Furlan & Hoshino, 2001; dePaula & Hoshino, 2002). Increased anxiety during the lactation period may also explain, at least partially, the results reported in the present paper. In this case, the significant increase in the number of longer episodes of awakenings observed in LRD may be interpreted as being the result of sleeplessness induced by an increased anxiety level. Postpartum insomnia seems to correlate to the emergence of psychiatric disorders (Karacan et al., 1968) and such considerations makes the LRD, forgotten in sleep research, a fascinating subject for investigation.

## REFERENCES

- Adels, L.E. and Leon, M. Thermal control of mother-young contact in Norway rats: factor mediating the chronic elevation of maternal temperature. *Physiol. Behav.*, 36: 183-196, 1986.

- Bates, A., Lori, A. and Leon, M. Thermal control of maternal contact bouts: the interbout interval. *Physiol. Behav.*, 34:835-837, 1985.
- Dement, W. The effect of dream deprivation. *Science*, 131:1705-1707, 1960.
- dePaula, H.M.G. Envolvimento de mecanismos convulsivos na etiologia das manifestações de pânico da defesa anti-predatória. Doctoral thesis. Universidade Estadual Paulista, Botucatu, 1999.
- dePaula, H.M.G. and Hoshino, K. Correlation between the fighting rates of REM sleep-deprived rats and susceptibility to the "wild running" of audiogenic seizures. *Brain Res.*, 926:80-85, 2002.
- Furlan, F.A. and Hoshino, K. Fighting by sleep-deprived rats as a possible manifestation of panic: effects of sodium lactate. *Braz. J. Med. Biol. Res.*, 34:359-366, 2001.
- Hoshino, K. and Toloi, J.R.G. Neocortical spindling during wakefulness in the rat. *Braz. J. Med. Biol. Res.*, 28:337-342, 1995.
- Hoshino, K. Food deprivation and hypothermia in desynchronized sleep-deprived rats. *Braz. J. Med. Biol. Res.*, 29:41-46, 1996.
- Johansen L., Toth, L., Rosenthal, R.S., Opp, MR., Obal Jr, F., Cady, A.B. and Krueger, J.N. Somnogenic, pyrogenic, and hematologic effects of bacterial peptidoglycan. *Am. J. Physiol.*, 260:R126-133, 1990.
- Jouvet, M. Neurophysiology of the states of sleep. *Physiol. Rev.*, 47:117-177, 1967.
- Karacan, I., Heine, W., Agnew Jr, H., Williams, R.L., Webb, W.B. and Ross, J.J. Characteristics of sleep patterns during late pregnancy and the postpartum periods. *Am. J. Obst. Gyn.*, 101:579-586, 1968.
- Kimura, M., Zhang, S.Q. and Inoué, S. Pregnancy-associated sleep changes in the rat. *Am. J. Physiol.*, 271: R1063-R1069, 1996.
- Knecht E.A., Toraason, M.A. and Wright, G.L. Thermoregulatory ability of female rats during pregnancy and lactation. *Am. J. Physiol.*, 239:R470-475, 1980.
- Krueger, J.N. Effects of cytokines on sleep patterns. *Physiologist*, 37: A-9, 1994.
- Landis, C.A., Bergman, B.N., Ismail, M.M. and Rechtschaffen, A. Sleep deprivation in the rat: XV. Ambient temperature choice in paradoxical sleep-deprived rats. *Sleep*, 15:13-20, 1992.
- Lenneberg-Hoshino, C. Mecanismos de equilíbrio populacional: avaliação do papel de uma atividade eletrocortical anormal do envelhecimento no processo de renovação das gerações de roedores. Doctoral thesis. Instituto de Biociências, Universidade Estadual Paulista, Botucatu, 1999.

- León, M., Groskerry, P.G. and Smith, G.K. Thermal control of mother-young contact in rats. *Physiol. Behav.*, 21:793-811, 1978.
- Lincoln, D.W., Hentzen, K., Hin, T., van der Shoot, P., Clarke, G. and Summerlee, A.J. Sleep: a prerequisite for reflex milk ejection in the rat. *Exp. Brain Res.*, 38:151-162, 1980.
- Moldofsky, H. Sleep and the immune system. *Int. J. Immunopharmacol.*, 17:649-654, 1995.
- Sandrin, M.F.N. Sono de ratos confinados em alta densidade populacional. Dissertação. Instituto de Biociências, Botucatu, 1996.
- Schmidek, W.R., Hoshino, K., Schmidek, M. and Timo-laria, C. Influence of environmental temperature on the sleep-wakefulness cycle in the rat. *Physiol. Behav.*, 8:363-371, 1972.
- Seabra, M.L. and Tufik, S. Sodium diclofenac inhibits hyperthermia induced by paradoxical sleep deprivation: the possible participation of prostaglandins. *Physiol. Behav.*, 54:923-926, 1993.
- Shinkoda, H., Matsumoto, K. and Park, Y.M. Changes in sleep-wake cycle during the period from late pregnancy to puerperium identified through wrist actigraphy and sleep log. *Psychiatry Clin. Neurosci.*, 53: 133-134, 1999.
- Timo-laria, C., Negrão, N., Schmidek, W.R., Hoshino, K., Menezes, C.E.L. and Rocha, T.L. Phases and states of sleep in the rat. *Physiol. Behav.*, 5:1057-1062, 1970.
- Voloschin, C.M. and Tramezzani, J.H. Milk ejection reflex linked to slow wave sleep in nursing rat. *Endocrinology*, 105:1202-1107, 1979.
- Voloschin, C.M. and Tramezzani, J.H. Relationship of prolactin release in lactating rats to milk ejection, sleep state, and ultrasonic vocalization by the pups. *Endocrinology*, 114:618-623, 1984.
- Yamaoka, S. Participation of limbic-hypothalamic structures in circadian rhythm of slow wave sleep and paradoxical sleep in the rat. *Brain Res.*, 151:255-268, 1978.
- Yamaoka, S. Modification of circadian sleep rhythms by gonadal steroids and the neural mechanisms involved. *Brain Res.*, 185:385-398, 1980.